





TABLE 1. Hyades *Caby* Photometry

ID	HIC/HD	Note	V	σ_V	$b - y$	σ_{by}	hk	σ_{hk}	N_{obs}	V_{avg}	$b - y_{avg}$	N_V, N_{by}	N_{sur}
1	15304	B	7.388	0.005	0.364	0.002	0.690	0.003	5	7.388	0.364	5	1
2	15310	B	7.774	0.008	0.394	0.006	0.727	0.007	6	7.773	0.396	9	3
4	16529		8.873	0.013	0.505	0.009	0.977	0.001	4	8.873	0.505	4	1
5	16908	B	9.332	0.002	0.533	0.009	1.086	0.014	3	9.332	0.533	3	1
7	18327		8.969	0.019	0.523	0.004	1.027	0.013	7	8.976	0.523	10	2
9	18719	NM	8.673	0.025	0.439	0.009	0.769	0.016	3	8.673	0.439	3	1
10	19148		7.839	0.013	0.381	0.013	0.676	0.010	3	7.842	0.384	9,14	6
13	19504		6.598	0.011	0.270	0.009	0.529	0.010	1	6.602	0.281	3,6	4
15	19793		8.040	0.014	0.421	0.006	0.733	0.010	4	8.042	0.420	6	3
16	19789		7.061	0.011	0.282	0.009	0.531	0.010	1	7.060	0.283	3,6	4
17	19781	B	8.459	0.010	0.436	0.007	0.774	0.009	13	8.458	0.434	19	3
18	19786		8.037	0.015	0.412	0.005	0.723	0.004	11	8.039	0.410	15	3
19	19796		7.108	0.007	0.335	0.006	0.599	0.007	5	7.104	0.338	12,16	6
21	19934		9.149	0.018	0.491	0.004	0.916	0.001	4	9.142	0.487	7	2
22	20019	B	8.320	0.013	0.467	0.008	0.809	0.001	5	8.317	0.466	10	3
23	20056	B	7.533	0.010	0.427	0.009	0.755	0.007	4	7.539	0.426	12	4
25	20082		9.559	0.012	0.558	0.005	1.188	0.011	6	9.561	0.555	14	3
26	20130		8.616	0.021	0.455	0.001	0.840	0.007	4	8.611	0.455	5	2
27	20146		8.472	0.013	0.447	0.002	0.820	0.008	6	8.469	0.444	13,15	5
29	20215	B	6.869	0.010	0.354	0.004	0.634	0.013	5	6.862	0.357	33,37	6
30	20219	B	5.587	0.008	0.171	0.005	0.467	0.006	14	5.586	0.175	15,19	3
31	20237		7.458	0.006	0.366	0.006	0.635	0.003	4	7.458	0.367	7,10	5
37	20357		6.607	0.006	0.274	0.003	0.494	0.006	16	6.604	0.276	19,24	5
39	20441	B	7.840	0.011	0.417	0.008	0.756	0.009	2	7.849	0.420	7	3
40	20440	B	6.982	0.015	0.367	0.002	0.665	0.017	3	6.978	0.370	6,9	4
42	20480		8.833	0.007	0.455	0.007	0.853	0.012	3	8.833	0.457	7	3
43	20482	B	9.386	0.021	0.529	0.006	1.022	0.009	3	9.386	0.529	3	1
46	20492		9.111	0.007	0.502	0.006	0.974	0.009	15	9.107	0.502	26	4
48	20557		7.136	0.011	0.347	0.009	0.597	0.013	2	7.132	0.346	8,12	6
49	HD 27835		8.220	0.009	0.382	0.002	0.701	0.004	7	8.222	0.382	12,17	4
50	20553	B	7.586	0.011	0.397	0.006	0.671	0.011	6	7.576	0.392	62,65	7
51	20567		6.984	0.010	0.300	0.001	0.551	0.001	2	6.969	0.300	7,11	5
52	20577	B	7.800	0.003	0.390	0.005	0.670	0.014	12	7.798	0.390	18,23	6
57	20661	B	6.459	0.006	0.320	0.007	0.575	0.004	13	6.448	0.323	17,21	3
58	20686	B	7.512	0.013	0.431	0.010	0.766	0.009	4	7.516	0.430	7	3
59	HD 28034	B	7.475	0.008	0.359	0.003	0.650	0.007	16	7.473	0.358	63,70	7
62	20712	B	7.374	0.021	0.347	0.001	0.657	0.003	3	7.369	0.348	8,12	5
63	20719	B	8.040	0.012	0.414	0.003	0.720	0.013	11	8.040	0.411	21,22	6
64	20741		8.130	0.006	0.419	0.005	0.754	0.011	13	8.114	0.417	25,27	7
65	20815		7.415	0.009	0.353	0.007	0.628	0.007	14	7.417	0.352	21,27	6

TABLE 1. (continued)

ID	HIC/HD	Note	V	σ_V	$b - y$	σ_{by}	hk	σ_{hk}	N_{obs}	V_{avg}	$b - y_{avg}$	N_V, N_{by}	N_{sur}
66	20826		7.508	0.006	0.359	0.002	0.635	0.002	4	7.505	0.364	7,10	4
68	20873	B	5.917	0.005	0.209	0.005	0.450	0.009	14	5.916	0.212	15,19	3
69	20890	B	8.610	0.009	0.447	0.004	0.834	0.013	3	8.609	0.448	4	2
73	20899		7.841	0.011	0.384	0.002	0.700	0.004	6	7.840	0.387	13,15	5
75	20916	B	6.560	0.003	0.346	0.001	0.612	0.002	3	6.568	0.349	15,19	7
77	20935	B	7.025	0.012	0.323	0.003	0.573	0.009	4	7.024	0.329	10,13	5
78	20948	B	6.898	0.006	0.304	0.006	0.543	0.004	4	6.901	0.304	10,15	6
79	20951	B	8.935	0.010	0.488	0.002	0.947	0.005	4	8.939	0.487	15,17	5
81	21008	B	7.072	0.022	0.310	0.006	0.544	0.004	2	7.085	0.317	5,8	5
86	21066		7.027	0.001	0.311	0.003	0.552	0.005	4	7.029	0.314	10,13	5
87	21099		8.592	0.006	0.454	0.005	0.841	0.017	5	8.591	0.455	6	2
88	21112		7.778	0.013	0.354	0.001	0.633	0.008	8	7.773	0.355	13,16	5
90	21152		6.372	0.004	0.275	0.010	0.522	0.007	10	6.372	0.278	10,14	2
91	HD 28783	B	8.923	0.012	0.520	0.002	0.956	0.008	3	8.923	0.520	3	1
92	HD 28805		8.670	0.021	0.455	0.010	0.843	0.008	5	8.663	0.454	11	3
93	HD 28878		9.401	0.002	0.519	0.001	1.015	0.006	3	9.389	0.517	7	3
94	21267		6.625	0.010	0.289	0.008	0.501	0.009	1	6.620	0.290	4,7	4
96	21280	B	8.497	0.024	0.499	0.011	0.932	0.010	3	8.497	0.499	3	1
97	21317		7.923	0.005	0.395	0.008	0.710	0.010	3	7.919	0.396	8	3
99	HD 29159		9.396	0.011	0.506	0.012	0.991	0.023	3	9.382	0.506	7	3
102	21543	B	7.532	0.012	0.383	0.004	0.682	0.008	4	7.531	0.390	7,12	4
103	21588	B	5.797	0.004	0.193	0.005	0.456	0.006	12	5.791	0.195	17,20	4
105	21637		7.517	0.012	0.362	0.008	0.676	0.009	2	7.517	0.360	2,5	2
106	21654	B	7.967	0.006	0.414	0.005	0.749	0.006	5	7.968	0.412	9	3
109	21741		9.399	0.011	0.479	0.013	0.909	0.021	3	9.399	0.479	3	1
113	22221	B	7.246	0.003	0.359	0.001	0.642	0.003	4	7.242	0.365	7,11	4
114	22265	B	8.533	0.008	0.435	0.002	0.831	0.007	2	8.530	0.444	6	3
115	22350	B	9.065	0.015	0.498	0.004	0.965	0.003	3	9.065	0.498	3	1
116	22380		8.966	0.003	0.496	0.008	0.936	0.010	3	8.968	0.496	4	2
118	22422		7.741	0.002	0.367	0.004	0.682	0.005	4	7.734	0.372	9,12	5
119	22496	B	7.089	0.006	0.360	0.004	0.623	0.016	3	7.094	0.364	5,8	4
120	22505	B	7.721	0.006	0.446	0.001	0.819	0.003	6	7.713	0.448	10	2
121	22524	B	7.294	0.009	0.323	0.002	0.600	0.005	6	7.288	0.329	13,16	5
122	22550	B	6.746	0.004	0.348	0.003	0.622	0.009	4	6.740	0.355	11,14	5
124	22607	B	6.252	0.010	0.319	0.001	0.551	0.005	3	6.245	0.328	10,15	5
127	23069		8.892	0.004	0.447	0.005	0.846	0.004	5	8.888	0.447	8	3
140	20601	B	8.928	0.011	0.465	0.001	0.845	0.001	4	8.927	0.465	9	3
142	22203	B	8.314	0.009	0.419	0.003	0.753	0.002	5	8.313	0.418	9	3
143	22566		7.895	0.006	0.339	0.008	0.624	0.010	4	7.896	0.342	7	3
162	19870	B	7.823	0.013	0.435	0.007	0.798	0.003	4	7.826	0.436	7	3

TABLE 1. (continued)

ID	HIC/HD	Note	V	σ_V	$b - y$	σ_{by}	hk	σ_{hk}	N_{obs}	V_{avg}	$b - y_{avg}$	N_V, N_{by}	N_{sur}
170	18018	B	10.176	0.010	0.670	0.008	1.443	0.021	2	10.176	0.670	2	1
172	HD 285762	NM	10.203	0.010	0.401	0.008	0.724	0.009	1	10.178	0.404	5	2
174	20563		9.975	0.017	0.602	0.005	1.314	0.020	3	9.984	0.600	10	3
177	HD 285828	B	10.281	0.020	0.613	0.011	1.335	0.046	2	10.300	0.619	8	3
178	20850		9.024	0.013	0.496	0.001	0.945	0.001	4	9.024	0.500	14	4
179	20827	B	9.480	0.008	0.537	0.004	1.097	0.009	6	9.475	0.536	9	2
180	20978		9.079	0.004	0.502	0.003	0.984	0.007	2	9.077	0.503	6	3
181	HD 285805		10.308	0.010	0.647	0.008	1.441	0.015	3	10.305	0.655	9	2
182	HD 28545	B	8.916	0.012	0.506	0.009	0.917	0.014	3	8.909	0.502	7	3
183	HD 28977		9.671	0.011	0.527	0.005	1.060	0.014	3	9.665	0.531	10	3
185	21762	B	9.430	0.005	0.621	0.008	1.253	0.018	3	9.434	0.627	6	2
187	23498	B	8.978	0.005	0.458	0.004	0.870	0.005	6	8.981	0.458	11	3
1068	19808		10.683	0.009	0.679	0.002	1.747	0.007	2	10.684	0.694	6	2
1135	20086	B	9.978	0.010	0.628	0.009	1.348	0.015	2	9.972	0.626	5	2
1407	20762		10.456	0.010	0.653	0.008	1.517	0.009	1	10.462	0.661	3	2
1627	21123	B	9.526	0.016	0.553	0.009	1.175	0.011	2	9.532	0.557	4	2
2010	19098		9.269	0.017	0.521	0.009	1.012	0.015	3	9.283	0.518	6	2
2011	18946		10.106	0.015	0.613	0.005	1.317	0.010	4	10.106	0.614	4	1
2014	19316		11.240	0.005	0.780	0.008	1.468	0.017	3	11.240	0.780	3	1
2015	19207		10.481	0.010	0.672	0.001	1.479	0.040	5	10.481	0.672	5	1
2016	19263		9.898	0.012	0.587	0.009	1.228	0.021	3	9.902	0.584	5	2
2018	HD 284785		9.430	0.014	0.533	0.005	1.077	0.002	4	9.430	0.533	4	1
2020	19591	B	9.385	0.008	0.643	0.007	1.077	0.007	3	9.385	0.643	3	1
2033	20419	B	9.774	0.004	0.672	0.005	1.504	0.031	4	9.775	0.672	6	2
2059	20751	B	9.434	0.002	0.599	0.001	1.213	0.006	3	9.429	0.596	6	2
2080	21723		9.990	0.016	0.605	0.006	1.363	0.028	3	9.999	0.606	5	2
2083		B	10.131	0.015	0.658	0.008	1.454	0.023	2	10.131	0.658	2	1
2090	HD 29896	B	9.815	0.007	0.570	0.007	1.215	0.017	3	9.820	0.571	6	2
2098	22654		10.268	0.002	0.616	0.002	1.341	0.002	3	10.268	0.616	3	1
2119	22177		10.873	0.016	0.725	0.013	1.525	0.036	3	10.873	0.725	3	1
7015	17766		10.805	0.017	0.781	0.018	1.569	0.029	2	10.805	0.781	2	1
HG 7-034	18322		10.103	0.019	0.619	0.009	1.338	0.021	3	10.103	0.619	3	1
HG 7-115	19441		10.053	0.012	0.683	0.005	1.526	0.027	4	10.053	0.683	4	1
8064	HD 284785		9.807	0.005	0.590	0.005	1.295	0.011	3	9.807	0.590	3	1








TABLE 2. Hyades Mean $hk, b - y$ Relation


$b - y$	hk	$b - y$	hk	$b - y$	hk
0.16	0.460	0.38	0.678	0.60	1.303
0.18	0.465	0.40	0.718	0.62	1.353
0.20	0.470	0.42	0.753	0.64	1.408
0.22	0.475	0.44	0.812	0.66	1.473
0.24	0.480	0.46	0.858	0.68	1.523
0.26	0.498	0.48	0.903	0.70	1.533
0.28	0.516	0.50	0.958	0.72	1.533
0.30	0.542	0.52	1.018	0.74	1.533
0.32	0.568	0.54	1.113	0.76	1.533
0.34	0.598	0.56	1.183	0.78	1.533
0.36	0.638	0.58	1.238	0.80	1.533

















Caby Photometry of the Hyades: Comparisons to the Field Stars

Barbara J. Anthony-Twarog, Bruce A. Twarog, and Judy Yu

Department of Physics and Astronomy, University of Kansas, Lawrence, KS 66045-2151

Electronic mail: bjat@ukans.edu, twarog@ukans.edu, judyyu@ku.edu

ABSTRACT

Intermediate-band photometry of the Hyades cluster on the *Caby* system is presented for dwarf stars ranging from spectral type A through late K. A mean hk , $b - y$ relation is constructed using only single stars without anomalous atmospheres and compared to the field stars of the solar neighborhood. For the F dwarfs, the Hyades relation defines an approximate *lower* bound in the two-color diagram, consistent with an $[\text{Fe}/\text{H}]$ between +0.10 and +0.15. These index-color diagrams follow the common convention of presenting stars with highest abundance at the bottom of the plot although the index values for the metal-rich stars are numerically larger. For field F dwarfs in the range $[\text{Fe}/\text{H}]$ between +0.4 and -1.0 , $[\text{Fe}/\text{H}] = -5.6 \delta hk + 0.125$, with no evidence for a color dependence in the slope. For the G and K dwarfs, the Hyades mean relation crosses the field star distribution in the two-color diagram, defining an approximate *upper* bound for the local disk stars. Stars found above the Hyades stars fall in at least one of three categories: $[\text{Fe}/\text{H}]$ below -0.7 , $[\text{Fe}/\text{H}]$ above that of the Hyades, or chromospherically active. It is concluded that, contrary to the predictions of model atmospheres, the hk index for cool dwarfs at a given color hits a maximum value for stars below solar composition and, with increasing $[\text{Fe}/\text{H}]$ above some critical value, declines. This trend is consistent, however, with the predictions from synthetic indices based upon much narrower Ca filters where the crossover is caused by the metallicity sensitivity of $b - y$.

Subject headings: open clusters:individual (Hyades)—stars:abundances—techniques:photometric

1. INTRODUCTION

Among well-studied open clusters, the Hyades occupies a unique niche due to the combination of its proximity and its high metallicity (Twarog et al. 1997). This rich sample of nearby stars of common age and composition provides a matchless testbed for any investigation of purely temperature-dependent trends at a given $[\text{Fe}/\text{H}]$ among dwarfs with a wide range of mass. In particular, the high metallicity of the Hyades has led to its adoption as a reference point for zeroing and/or testing the metallicity scale for several photometric systems, including the *UBV* (Carney 1979), the *DDO* (Deming et al. 1977; Twarog & Anthony-Twarog 1996), and the Strömgren (Crawford 1975; Schuster & Nissen 1989) systems. The purpose of this investigation is to add another, the *Caby* system, to the long list of fun-

damental color relations defined by the Hyades, while investigating the effects of metallicity, and potentially of age, on the hk index for a large sample of nearby field stars.

The *Caby* system represents an extension of the traditional four-color, *uvby* intermediate-band photometric system to a fifth filter centered on the H and K lines of Ca II. Details of the filter definition and design as well as the fundamental standards may be found in Anthony-Twarog et al. (1991), while an extensive catalog of stars observed on the system and tied to the $b - y$ scale of Olsen (1993) may be found in Twarog & Anthony-Twarog (1995). The filter was designed initially with metal-deficient stars in mind, as demonstrated by numerous applications to date on normal field stars (Anthony-Twarog et al. 1991, 1992; Anthony-Twarog & Twarog 1998; Anthony-

Twarog et al. 2000), clusters (Anthony-Twarog et al. 1995; Rey et al. 2000), and variables (Baird 1996; Hintz et al. 1998). Metallicity calibrations have been produced for both the metal-deficient giants (Anthony-Twarog & Twarog 1998) and metal-deficient dwarfs (Anthony-Twarog et al. 2000), but preliminary analysis indicated that for dwarfs hotter than the sun, the hk index, defined as $(Ca-b)-(b-y)$, remains metallicity sensitive for stars of solar abundance or higher (Twarog & Anthony-Twarog 1995), a result consistent with the theoretical models of Soon et al. (1993). Because of the high metallicity of the Hyades relative to the typical star in the field of the solar neighborhood, it provides an ideal test of this prediction, as well as a means of probing the limits of its sensitivity.

A second issue of particular relevance for metal-rich stars is the role of chromospheric emission on the apparent strength of the hk index. As one moves toward lower temperature, the expectation is that the absorption features will saturate, leaving an index which is almost exclusively dependent upon temperature. That the sensitivity of the hk index to metallicity should decline near $[Fe/H] = 0.0$ for $(b-y)$ redder than 0.5 is readily apparent in the synthetic indices of Soon et al. (1993), though the models imply that even G and K dwarfs with abundances well above solar should exhibit larger hk indices at a given color. However, line reversals triggered by chromospheric activity (see, e.g., Vaughan 1980) may fill in the cores of the absorption features, making the star appear more metal-deficient than it actually is. Moreover, the strength of this line reversal has long been known to be age-dependent (Wilson 1963). Since the Hyades is moderately young (less than 1 Gyr) compared to the average cool dwarf near the sun and contains some stars with an anomalous degree of chromospheric activity, it may provide some insight into this question. Finally, the majority of the stars known to be Hyades members have been studied in detail, thereby allowing us to look for additional photometric anomalies not tied to the metallicity of the stars.

Section 2 contains the details of the observations, their transformation to the standard *Caby* system, and the merger of our V , $b-y$ data with an extensive array of published data on the Hyades dwarfs in an effort to minimize the potential ef-

fects of internal photometric scatter on the mean relations. In Sec. 3 we derive the Hyades mean relation in the hk , $(b-y)$ diagram and discuss the potential sources of intrinsic scatter in the two-color diagram. Sec. 4 compares the single-star Hyades relation to a large sample of nearby field stars, providing some insight into possible sources of the differences between expectation and reality, particularly at cooler temperatures. Sec. 5 contains a summary of our conclusions and suggestions for further work on the system in light of the Hyades anomalies.

2. The Data

2.1. Observations

The data discussed here were obtained as part of the wide-ranging program of observations on the *Caby* system started in Dec. 1983 and continued on through 1994. The *Caby* work since then has focused on more specialized applications, but some Hyades stars were included from the beginning of the program with calibration and standardization purposes in mind. The preliminary sample was selected from the well-known, brighter members identified by Van Bueren (1952), but the more comprehensive database detailed here is tied to the compilation of Griffin et al. (1988). The observations were made on a variety of telescopes including the 0.9 m and 1.2 m at KPNO and, primarily, the 0.6 m, 0.9 m, 1.0 m, and 1.5 m at CTIO. All the telescopes at the national observatories were equipped with pulse-counting, single-channel photometers mounted with S-20 photomultipliers. In addition, a modest number of observations were obtained using a pulse-counting photometer equipped with a 1P21 photomultiplier on the 0.4 m telescope at Braeside Observatory in Arizona. The filter sets used have changed over the years, but extreme care has been taken to ensure transformation to the standard system and no significant problems have been encountered. Observations of the fainter stars in this study were all made with the 1.0 m and 1.5 m telescopes at CTIO equipped with the same filter set. For more information regarding the design of the *Ca* filter and the observing procedure, the reader is referred to Anthony-Twarog et al. (1991).

2.2. Reduction and Transformation

A complete description of the methodology used to merge all the observations taken from the multiple runs over a number of years is given in Twarog & Anthony-Twarog (1995) and will not be repeated here. A key point is that the definition of the standard $b - y$ system was changed between the publication of the primary standards (Anthony-Twarog et al. 1991) and the final catalog. This was prompted by the extension of the system to include a significant fraction of cool dwarfs and the publication of the definitive set of cool star observations on the $uvby$ system by Olsen (1993). The composite, homogenized sample on the instrumental system included all of our Hyades data up to that time; this sample was used to define the transformation in $b - y$, with separate transformation curves defined for hotter stars, cool dwarfs, and cool giants. In the mean, the large overlap between the *Caby* and $uvby$ catalogs produced an excellent match between the two systems. Though several brighter stars in the Hyades were included in the 1995 catalog, the fainter dwarfs were not due to the optimistic expectation of additional observations.

We have accumulated over 535 observations on the system of the *Caby* catalog (Twarog & Anthony-Twarog 1995) for 114 Hyades dwarfs. As the $b - y$ colors of that catalog were intended to be consistent with Olsen (1993), the Hyades data should be also; indeed, a few of the brighter Hyades stars were included in the analysis from which the color transformations to Olsen (1993) were developed. Yet, it is not uncommon for cluster photometric samples to show small offsets with respect to a standard system even though the all-sky data reveal no systematic differences. We were particularly concerned because most of the Hyades stars were observed at CTIO at large airmass. To minimize the discrepancies with the Olsen (1993) system, we directly compared the Hyades stars common to both catalogs; Hyades stars from Olsen (1994) were also included in this comparison.

The results of the comparison revealed slight differences in our colors and magnitudes with respect to Olsen (1993). For 81 stars bluer than $b - y = 0.50$, our preliminary colors were too red by 0.0049 ± 0.0044 (s.d.). The V magnitudes are correspondingly brighter by 0.006 ± 0.018

(s.d.). For the 26 redder stars, a linear relation between the $b - y$ colors was derived; the transformation between our preliminary colors and those of Olsen (1993) was $(b - y)_{OLS} = 1.049(b - y)_{ATT} - 0.0264$. The points scatter about the mean relation with residuals amounting to only ± 0.0055 mag (s.d.). A similar adjustment was required for our preliminary V magnitudes, i.e., $V_{OLS} = V_{ATT} + 0.1(b - y)_{ATT} - 0.0391$ with a scatter of ± 0.0086 . The new Hyades *Caby* data presented in Table 1 (columns 4 through 9) have been adjusted from our preliminary results according to these precepts and are solidly tied to the Olsen (1993) system. The identification number found in the WEBDA database is found in column 1, with HIC, HD or other identifications listed in column 2. Notes in column 3 refer to information gathered about suspected binaries and non-members, described further in Sec. 3.

In an effort to further improve the final precision of the Hyades $b - y$ colors and magnitudes, we searched the internet open cluster database, WEBDA, to find Strömgren photometric datasets in the Hyades which substantially overlapped with our Hyades data. Published $V, b - y$ data from Olsen (1983, 1993, 1994), Reglero et al. (1992), and Sowell & Wilson (1993) were used. Most of the other sizable Hyades Strömgren photometric samples are restricted to brighter and hotter main sequence stars and were not included in the merger. Of the datasets included in the merged sample, that of Crawford & Perry (1966) does not include V magnitudes. One large sample, that of Perry (1969) provided only a rather noisy comparison to our $(b - y)$ colors and was not used. Finally, we also included unpublished data for 10 Hyades dwarfs obtained by Bruce Carney with the original prototype of the Ca filter; only $(b - y)$ colors and hk indices were available.

As expected, small offsets between these surveys and our preliminary sample were found, though with no evidence for color terms. Typical differences in $b - y$ ranged from -0.005 to $+0.015$ mag, with a range of ± 0.016 for average differences in V magnitudes. The indices from each dataset were adjusted for these small differences, placing them on our preliminary system, then transformed to the Olsen (1993) system according to the precepts indicated above. The data sets were finally merged and averaged using

weights based upon the consistency between each dataset and our preliminary Hyades dataset. The final averaged V magnitudes and $b - y$ colors are found in columns 11 and 12 of Table 1, followed by the number of V and $b - y$ observations incorporated in the average and the number of datasets included. The standard errors of the mean for the final magnitudes and indices were quite small and are not included in Table 1. Typical values of ± 0.004 (s.e.m.) and ± 0.003 (s.e.m.) were found for V and $b - y$, respectively.

Finally, the $b - y$ colors for Hyades stars in common with Olsen (1993) were checked to verify that the color system is consistent with that sample. From 81 stars in common, the mean difference in $(b - y)$ for the merged sample is -0.0010 ± 0.0037 (s.d.) in the sense (OLS - Table 1).

3. The Hyades $hk - (b - y)$ Relation

In defining any two-color relation, one ideally wants to exclude stars affected by anomalies which might distort the relation beyond normal photometric scatter. In the case of a cluster sample, a fundamental requirement is that stars be classified as cluster members. Since the time of the original compilation of the Hyades stars for observation, based primarily upon the discussion in Griffin et al. (1988), additional parallax, radial-velocity, and proper-motion observations, particularly those by *Hipparcos* (Perryman et al. 1998), have clarified and eliminated the membership of a few stars in Table 1. Star VB/WEBDA 9, despite positions in the color-magnitude diagram (CMD) and $hk, b - y$ diagram consistent with the Hyades, is now classed as a kinematic non-member. A second, more obvious interloper is VB/WEBDA 172. Originally classed as a Hyades subdwarf, it lies well below the main sequence in the Hyades CMD. This class of stars is now known to be a product of more distant nonmembers whose space velocities conspire to produce proper motions compatible with those of the Hyades members (Hanson & Vasilevskis 1983). These two stars will be removed from further discussion. The remaining 112 stars are classed as probable members by Perryman et al. (1998) or, if not included in the *Hipparcos* survey, have been tagged as probable members by Griffin et al. (1988).

The next obvious exclusion to make is the elimi-

nation of all stars which are multiple systems. The presence of one or more companions may distort the colors of the composite system directly via the combined colors of the stars or indirectly through an alteration of the structure and/or evolutionary state of the star and its atmosphere, as in the case of chromospherically active stars created through tidal interaction. Following de Bruijne et al. (2001), in deriving the mean relation we have eliminated any star which has been tagged as a potential binary through direct observation from the ground or from data obtained from *Hipparcos*; see Table A.1 of de Bruijne et al. (2001). These classifications have been supplemented by the regularly updated information on the Hyades from WEBDA, in particular the interferometric survey of Hyades multiplicity by Patience et al. (1998). In many cases, the binary nature of the system may be expected to have little or no impact on the observed colors; the angular separation of the stars might be too large for both stars to be observed simultaneously with a standard photometer or the mass, temperature, and luminosity ratio between the stars may be too large to affect the colors of the primary star. With a reasonably extensive photometric sample, we have chosen to be very conservative in selecting stars to define the mean relations which can be definitively classed as single. Of the 112 members in Table 1, 54 stars have some indication of multiplicity and are noted with a “B” in column 3, leaving 58 single stars.

Before compiling the final relation, as a precaution in case some binaries have been included in the derivation due to inadequate information, one can construct artificial binary systems and estimate what impact they might have on the two-color relation. Use has been made of the final relation in constructing the composite pairs by choosing a star of a given color on the main sequence, then sequentially combining the fluxes in each bandpass for a series of stars at decreasing mass/luminosity along the main sequence. The locations of the composite systems in the CMD are illustrated in Fig. 1a, while the composite colors produce the pattern shown in Fig. 1b. In Fig. 1a, the curves at each location along the CMD start with two stars of identical color, then illustrate the effect of altering the composite system using a secondary which resembles the next sequential point down on the main sequence. The trend in

the CMD is well-known and essentially the same for all photometric systems. Composites of identical stars are located at the same color as the single stars, but 0.75 mag above the main sequence. As the secondary star is replaced with a cooler, less luminous star farther down the main sequence, the color difference initially shifts the composite increasingly to the red but at a smaller distance above the main sequence. Eventually, the decreasing luminosity of the secondary creates a smaller impact on the primary system; the color and absolute magnitude of the composite move closer to that of the primary star alone.

For the two-color diagram, the pattern of changes is slightly more complex because of the changing slope of the mean relation. For the stars hotter than the sun ($b - y \leq 0.4$), the composite systems are always redder in $b - y$ than the single stars, with the size of the shift larger for composites with the smaller difference in luminosity. The combined change in both indices moves the stars approximately parallel to the mean relation, implying that binaries should not be a significant source of scatter among hotter dwarfs in the $hk, b - y$ relation. The steepening slope for the late G through K dwarfs adds a twist to this pattern comparable to what is seen in the CMD. The composite system is always redder than the single primary, but the pattern in $b - y$ is to grow redder initially as the secondary is shifted down the main sequence, hit a maximum difference, then shift blueward toward the color of the primary star. Moreover, the combined effect of changing $b - y$ and hk places the composite star increasingly above the mean relation before finally approaching the values of the single primary. We conclude then that if composite systems are present within the sample, they should have little to no impact among the hotter stars. Among the K dwarfs, companions should shift the system above the mean relation, *i.e.*, toward smaller hk , with the size of the shift depending upon the exact colors of the two stars. Maximal shifts between 0.05 and 0.10 mag are expected in the $b - y$ range of 0.45 to 0.65.

Using the 58 single Hyades members, a preliminary two-color relation was derived by sorting the sample in $b - y$ and, starting at the blue end, constructing a five-point, weighted average which then shifted redward by one point before repeating

the process. This was continued until a running average was available over the entire color range of the sample. This mean relation was checked for deviant points whose inclusion produced significant changes in slope over small changes in $b - y$; one point was removed. The averaging process was repeated, producing a mean relation which showed a smooth and continuous change in slope over the entire color range. Finally, residuals were calculated by comparing the data points to the mean relation and tested to ensure that the scatter about the mean curve averaged 0.0 with no color dependence. The final Hyades $hk, b - y$ relation is listed in Table 2 and shown as the solid line in Fig. 2. The open circles of Fig. 2 are the 57 single, non-deviant stars used to define the relation, while the crosses are the stars tagged as possible binaries. The two weakest parts of the relation are the extreme ends of the color range. For $b - y$ below 0.27 and greater than 0.7, the number of points available to define the curve is small and may be affected by possible binaries. At the cool end, the small numbers were set by the apparent faintness of the stars while at the blue end, hotter stars were too bright ($V \leq 5.5$) for even the smallest telescope accessible to this project. In both of these color ranges, it has been assumed that the Hyades relation has the same relative shape as that found for the field stars of comparable color. The Hyades members have been used essentially as a means of fixing the location of the curve in these regions, not defining the shape of the curve. In short, the extreme ends of the relation should only be used with caution.

Excluding the one deviant star, the 57 single stars scatter about the mean two-color relation with average residuals in hk , in the sense (OBS-REL), of $+0.001 \pm 0.019$ mag (s.d.). If binary stars are considered together with single stars, and if two additional deviant stars are excluded, the remaining 109 stars show average residuals of -0.003 ± 0.022 , indicating, as predicted, that binarity moves a star toward lower observed hk relative to the mean relation. The three stars which exhibit the largest deviation from the mean relation are the binary stars HIC 19591 and VB/WEBDA 185 (filled squares in Fig. 2), while the deviant single star is HIC 19808 (HG7-135; filled circle in Fig. 2). The single star is peculiar in that its colors imply that the absorption in the region of Ca II

H and K is anomalously large. Since the two-color relation flattens out in the $b-y$ region occupied by this star (~ 0.7), one cannot construct the indices demonstrated by this star through combinations of any stars on the standard relation. Moreover, as will become apparent in the next section, no field star has been observed which occupies a comparable region of the $hk, b-y$ diagram. Since one would expect that most forms of stellar interaction would enhance chromospheric emission leading to line filling, the source of the anomaly remains a mystery. The simplest solution may be that the observations are in error, though the star was observed twice and the agreement between the observations was quite good.

In contrast, the position of HIC 19591 (BD +23° 635), over 0.3 mag above the mean relation is readily explained. It is known to be a short-period, tidally-locked triple system that exhibits high levels of chromospheric activity (Griffin & Gunn 1981; Bopp et al. 1986) and an anomalous Li abundance (Thorburn et al. 1993). VB/WEBDA 185 is a composite composed of a K3 and a K7 dwarf, separated in absolute magnitude by only 1.6 mag (Griffin & Gunn 1981). The system does not show significant evidence for anomalous chromospheric emission (Stauffer et al. 1991). The deviation from the mean relation reflects the optimum difference in color and intrinsic luminosity required to shift the location of the system relative to that of the mean relation. The intrinsic colors of the stars place them in the ideal $b-y$ regime for shifts between 0.05 and 0.10 mag in hk relative to the mean relation, as observed.

4. The Field Star Comparison

With the Hyades $hk, b-y$ relation established, the next task is to determine what, if anything, distinguishes the Hyades relation from the typical field star in the solar neighborhood. As stated earlier, expectations are that the Hyades will establish an approximate lower bound for hotter field stars due to its high $[\text{Fe}/\text{H}]$, but remain typical of most stars at lower temperatures due to the saturation of the Ca II lines. To ensure that any anomalies between the Hyades and the field stars are not a product of zero-point offsets within the photometry, one additional check has been made. In Sec. 2 it was demonstrated that our merged V

and $b-y$ data are consistent with the system of Olsen (1993). If we compare the $hk, b-y$ indices for Hyades stars to field dwarfs from the Twarog & Anthony-Twarog (1995) catalog, can we be certain that the dwarfs in that catalog are indeed on a color system consistent with Olsen (1993)?

A set of 401 dwarfs was compiled representing the overlap between the catalogs of Olsen (1993) and Twarog & Anthony-Twarog (1995). The stars cover a brightness range from $V = 4.8$ to $V = 11.6$, and a color range from $(b-y) = 0.08$ to 1.1; the great majority of these stars have colors between $(b-y)$ of 0.2 and 0.7. No giants were included in this overlap set, and a number of stars with discrepant V magnitudes were excluded. The mean differences in the sense (OLS93 - ATT95) are 0.0007 ± 0.0006 (s.e.m.) for V and -0.0007 ± 0.0003 (s.e.m.) for $(b-y)$. There are no discernible trends with respect to color or magnitude for ΔV or $\Delta(b-y)$, confirming the comparison made during the construction of the original catalog (Twarog & Anthony-Twarog 1995).

4.1. The Hotter Dwarfs

To illustrate the trends in the two-color diagram at reasonable resolution, we divided the sample in two color ranges with the break point at $b-y = 0.45$. For the hotter stars not classified as giants, Fig. 3 shows the field star sample from Twarog & Anthony-Twarog (1995) with the Hyades mean relation superposed as a solid line. Over the F-star spectral range, $b-y$ between 0.2 and 0.4, the mean relation defines an approximate lower bound to the field star sample. This is expected if the hk index is predominantly a metallicity indicator and the scatter of stars below the Hyades is, in fact, composed mostly of stars more metal rich than the Hyades. The few stars that fall well below the Hyades relation and below virtually all the field stars are supergiants and not relevant to the discussion.

We wished to probe the metallicity sensitivity of the hk index as well as investigate the possibility that it might saturate, or even reverse, as a metallicity indicator for $[\text{Fe}/\text{H}]$ values greater than the Hyades. One relatively straight-forward way to do this is to construct an index, δhk , that describes the difference, in the sense (Hyades - star), of a star's hk index from the Hyades $hk, b-y$ sequence, and then compare δhk to photometric or

spectroscopic determinations of $[\text{Fe}/\text{H}]$.

Overlap of the hk catalog (Twarog & Anthony-Twarog 1995) with catalogs of high dispersion spectroscopic abundances is not tremendously large, consisting of 50 non-binary stars from the catalogs of Edvardsson et al. (1993) and Chen et al. (2000) that fall within the $b - y$ limits of 0.22 and 0.38, a color range for which we may also utilize the photometric metallicity calibration for F stars of Schuster & Nissen (1989). Chen et al. (2000) demonstrate effectively that the $[\text{Fe}/\text{H}]$ values in these two spectroscopic catalogs are mutually consistent; Fig. 4 illustrates the relationship between $[\text{Fe}/\text{H}]_{\text{SPEC}}$ and δhk for 50 F dwarfs. The form of the relationship is $[\text{Fe}/\text{H}] = -5.60\delta hk + 0.065$, where the intercept implies the $[\text{Fe}/\text{H}]$ value for $\delta hk = 0.0$, *i.e.*, Hyades abundance. The dispersion of points about the mean line, 0.08 dex, makes this result formally consistent with a presumed abundance for the Hyades of $[\text{Fe}/\text{H}] = 0.125$.

We compared the performance of δhk relative to photometric metallicity indicators as well, a considerably less straightforward task but one that recovers the advantages of several large Strömgren catalogs. The F star calibration of Schuster & Nissen (1989) is ideal for this temperature range and is constrained by limits on applicable $b - y$, m_1 and c_1 . A few features deserve note: several Hyades stars were included in the development of the calibration, but it was not primarily intended for use at near-solar abundances, an issue that can be addressed by small additive corrections to the $[\text{Fe}/\text{H}]$ scale. Small differences that exist between large catalogs on the Strömgren system can lead to slightly different results as well. The Schuster & Nissen (1989) calibration is consistent with their photometry; other catalogs may be transformed to their catalog system using published precepts. For example, 37 Hyades F dwarfs are included in the photometric catalog of Olsen (1983). If the abundance calibration of Schuster & Nissen (1989) is used, an abundance $[\text{Fe}/\text{H}] = -0.01 \pm 0.09$ is obtained. A slightly different photometric catalog, Olsen (1993) also contains 20 Hyades F dwarfs. The photometric calibration provides an estimate of $[\text{Fe}/\text{H}] = -0.06 \pm 0.07$ for these indices. This estimate is raised if the Strömgren indices of the Olsen (1993) catalog are mapped to the system of the calibration, Schus-

ter & Nissen (1989). These precepts are provided by Olsen (1993) and produce an estimate of $[\text{Fe}/\text{H}]$ for the Hyades stars of 0.04 ± 0.09 , about 0.1 dex higher than untransformed indices imply. Thus, slight differences between photometric scales can lead to modest differences in calculated abundances which may be adjusted by zero point corrections. There are also concerns about the **scale** of the photometric estimators of $[\text{Fe}/\text{H}]$ for F stars. Alonso et al. (1996) suggest an adjustment to abundances determined using the Schuster & Nissen (1989) F star calibration of the form: $[\text{Fe}/\text{H}]_{\text{SPEC}} = 1.22[\text{Fe}/\text{H}]_{\text{PHOT}} + 0.125$, where the intercept is fixed to ensure a Hyades abundance of +0.125.

With these points in mind, we constructed a comparison of δhk to photometric abundances, using the two largest, high-precision, F-star catalogs on the $uvby$ system with the greatest overlap with the hk catalog (Twarog & Anthony-Twarog 1995), those of Grønbech & Olsen (1976) for the brightest stars and Olsen (1983) for stars at fainter magnitudes. For each star between $b - y = 0.22$ and 0.38 common to the $uvby$ and hk systems, δhk is calculated while $uvby$ data for each star was processed through the F-star metallicity calibration of Schuster & Nissen (1989), using the limits on the range of indices for which the calibration is applicable to further restrict the sample. If the indices from Grønbech & Olsen (1976) and Olsen (1983) are mutually consistent, stars of Hyades abundance should have δhk values of 0.0. Without the adjustment to slope or zero point, the photometric calibration for Hyades stars yielded a value of $[\text{Fe}/\text{H}] = 0.0$, which is increased to 0.125 by the linear adjusted scale.

With these assumptions in mind, Fig. 5 provides a comparison of photometric estimates of $[\text{Fe}/\text{H}]$ to δhk values for 324 stars. The circles represent stars from Olsen (1983), the crosses data from Grønbech & Olsen (1976). A number of conclusions can be derived from this diagram. First, there is no inconsistency of slope or zero point to indicate that the two photometric samples are anything other than mutually consistent. As hoped, at $\delta hk = 0.0$, *i.e.*, Hyades abundance by definition, the implied $[\text{Fe}/\text{H}]$ value lies between +0.05 and +0.15, demonstrating that the Hyades metallicity scale is consistent with that defined by the field stars. A line fitted through the points

has a slope of -5.8, an intercept of +0.10, and a dispersion of ± 0.12 dex, expected from photometric errors alone and in spite of no applied corrections for reddening or any inhomogeneities within the samples of stars. The value of the slope is close enough to that derived from data presented in Fig. 4 to validate the amplification of the photometric abundance scale advocated by Alonso et al. (1996). It is concluded that for F dwarfs, the Hyades mean relation defines a reliable isometallicity line at $[\text{Fe}/\text{H}] = +0.125$.

4.2. The Cooler Dwarfs

Fig. 6 shows the unevolved cooler field stars from Twarog & Anthony-Twarog (1995) relative to the Hyades mean relation. Surprisingly, the Hyades relation crosses over the color-color band defined by the field stars and becomes an approximate upper bound to the sample, rather than a lower limit. This pattern extends from $b - y \sim 0.5$ to 0.65, where the flattening of the Hyades relation, coupled with the small number of points defining the mean relation, makes it difficult to pin down exactly where the Hyades is located relative to the field stars.

If the peculiar location of the Hyades relation is not the product of photometric error, what might be its cause? Three possibilities come to mind:

1) *The location of the two-color relation is linked to the Hyades anomaly.* The Hyades anomaly refers to observation that the unevolved F stars in the Hyades have c_1 indices that place them above the unevolved field stars at the same temperature (Crawford & Barnes 1969). Since c_1 is an indicator of surface gravity through the Balmer discontinuity, the higher c_1 values at a given $b - y$ or $\text{H}\beta$ imply that the F dwarfs in the Hyades are more luminous/evolved than the field stars of comparable age. Over the last 15 years, numerous solutions have been proposed and questioned, including filter errors (Eggen 1994; Joner & Taylor 1995), starspots triggered by chromospheric activity (Rose 1984; Soderblom 1989), some form of abundance anomaly, including overall metallicity (Barry 1974; Alexander 1986), CN variations (Eggen 1982; Brown & Twarog 1983), and helium (Strömberg et al. 1982; Alexander 1986). Others have questioned the reality of the phenomenon, though the work of Dobson (1990) indicates that something that is not predicted by simple, stan-

dard models is happening. Finally, Nissen (1988) would appear to confirm that the Hyades cluster is not alone in this peculiarity, while demonstrating that the anomaly is not a simple age or metallicity effect.

There are two reasons to discount the relevance of the Hyades anomaly in the current discussion. First, if a star has a higher c_1 index at a given color, this indicates that the star is fainter than expected in the ultraviolet region of the spectrum. The cooler Hyades dwarfs look metal-deficient, *i.e.*, their energy distribution is skewed to higher ultraviolet flux, though one could claim that over the narrow bandpass of the *Ca* filter alone, the flux is higher than expected. Second, photoelectric observations of the cooler stars by Reglero et al. (1992) demonstrate that the Hyades dwarfs have c_1 indices consistent with the mean relation for field stars and show no evidence for the anomaly identified among the hotter stars.

2) *The stars in the Hyades are younger than the average field star, so the increased chromospheric activity compensates for the higher metallicity by filling in the *Ca II H* and *K* lines and making the star appear metal-deficient.* If correct, stars that lie above the Hyades relation in the two-color diagram must be metal-poor and/or chromospherically active, the latter due to a younger age or environmentally-induced activity. The fact that the observed stars known to be chromospherically active to an anomalous degree sit well above the Hyades relation is consistent with this option.

3) *Though the metallicity effect on the *Ca II* lines should increase the value of *hk* at a given $b - y$ as $[\text{Fe}/\text{H}]$ increases, above a critical value of $[\text{Fe}/\text{H}]$ the index hits a maximum and then decreases with increasing $[\text{Fe}/\text{H}]$.* Among the most metal-rich stars in the solar neighborhood, Hyades members are positioned **above** stars of lower metallicity in the two-color diagram. Therefore, stars even more metal rich than the Hyades should fall above the mean relation in Fig. 6.

To sort through the possibilities, we have identified 59 stars that lie above the Hyades mean relation between $b - y = 0.50$ and 0.8. As a first step, the *Hipparcos* catalog was searched and all but 5 stars had a measured parallax. Of the remaining 54, all but 8 had σ/π of 0.16 or less, implying that reliable absolute magnitudes were possible. The CMD for the remaining 46 stars is

shown in Fig. 7 where the error bars drawn are based upon the one sigma uncertainty in the parallax. What is immediately apparent is that the sample contains a number of stars that should legitimately be classified as subgiants/giants, *i.e.*, any star with M_V below 5.0. To this sample of 8 stars can be added 3 stars that have measured parallaxes with reasonable errors, but absolute parallaxes so small the stars must be at large distance and therefore must be intrinsically more luminous than a dwarf. The misclassification of these stars as dwarfs explains their anomalous location in the two-color diagram because the mean relation for solar-metallicity giants is shifted approximately 0.1 mag above that for the dwarfs due to surface gravity effects. Though these stars will be excluded from further discussion, we note that two of the stars are of interest because they are anomalous even for giants. HD 13435 is the reddest star at the base of the giant branch, occupying a position that is undoubtedly heavily dependent upon its exceptionally high $[\text{Fe}/\text{H}]$ (Favata et al. 1997). HD 89499 sits well above the mean relation due to classification as both a subgiant and a chromospherically-active, short-period, tidally-locked binary (Ryan & Deliyannis 1995).

The next step in sorting through the remaining 48 stars is to estimate their metallicity and determine if they fall into either extreme end of the abundance range or cover the entire spread. As one might expect, the information available for the stars is mixed. The spectroscopic abundance catalog of Cayrel de Strobel et al. (2001) includes high dispersion results for 10 of the stars, 11 if one includes the value for HD 61606A as representative of HD 61606B. For stars with multiple abundance estimates, an unweighted average has been taken. Beyond spectroscopy, we can appeal to photometry for abundances. A survey of the *uvby* catalog of Hauck & Mermilliod (1998) identifies 18 additional stars with a full set of *uvby* indices, of which 11 fall within the calibration limits of the G-star calibration of Schuster & Nissen (1989). For stars with $[\text{Fe}/\text{H}]$ below -0.4 , the abundances derived from the calibration were adopted as is without any adjustment to the zero-points of the photometry or the metallicity scale. For stars with $[\text{Fe}/\text{H}]$ above this limit, the photometry was also uncorrected for any zero-point differences with Schuster & Nissen (1988), but the photometric abundances

were derived by comparison to the photometric abundance derived for Hyades stars at the same color using the Schuster & Nissen (1989) calibration. The reason for this change is discussed in detail in Twarog et al. (2002) and arises from a flaw in the photometric abundance calibration that leads to a severe underestimate of $[\text{Fe}/\text{H}]$ for cooler G and early K dwarfs at solar abundance and above. For derivation of the final abundances, it has been assumed that $[\text{Fe}/\text{H}]$ for the Hyades is $+0.125$. Finally, Alonso et al. (1996) have derived a photometric abundance of -1.5 for HD 4967, while Laird et al. (1988) obtain $+0.32$ for HD 95741.

Of the 24 stars with some semblance of an abundance estimate, 9 were below $[\text{Fe}/\text{H}] = -0.70$, implying that their position in the two-color diagram is the product of low $[\text{Fe}/\text{H}]$. Of the remaining 15 stars, none were found between $[\text{Fe}/\text{H}] = -0.20$ and -0.70 . The abundance distribution for these stars is plotted in Fig. 8 and clearly shows that, despite the lack of homogeneity in the sample abundances, the stars are heavily weighted toward Hyades abundance or higher, as predicted if the shift of the mean relation is due to a metallicity effect. All stars with $[\text{Fe}/\text{H}]$ below -0.70 are contained in the lowest bin.

The final classification property to check is that of chromospheric activity. Seven of the stars are notable for their unusual degree of chromospheric activity, in many cases leading to classification as *BY Dra* stars. HD 89499 is a subgiant and has been noted earlier. BD $-0^\circ 4234$ is a metal-poor dwarf in a tidally-locked binary system (Ryan & Deliyannis 1995) and was included in the metal-poor classification above. Thus, the presence of 5 additional stars above the Hyades in the two-color relation can now be clarified, leaving 19 with insufficient information to determine their status. An attempt has been made to discover if there is any general trend with chromospheric activity beyond the obvious effect for stars of extreme activity. The stellar sample with measurements of the chromospheric index, R'_{HD} compiled by Rocha-Pinto et al. (2000) has been matched with the *Caby* catalog, producing an overlap of 189 stars. Unfortunately, only 14 stars lie within the critical color range redder than $b - y = 0.50$. For each of these stars, a residual $\delta R'_{HK}$ was calculated by subtracting the observed index from a linear lower

limit drawn through the sample as a function of color. No correlation was found between this chromospheric index and the position of the star relative to the Hyades in the $hk, b - y$ diagram at the same color though, again, the sample is small.

We note that in some cases, stars have been categorized in a way that might indicate anomalously high $[\text{Fe}/\text{H}]$, *e.g.*, inclusion in a metal-rich moving group, but such a designation was not deemed reliable enough to justify the assumption of high $[\text{Fe}/\text{H}]$. Moreover, among the coolest dwarfs, many were selected because of their specification as being within such a group, making the sample less than random.

In Fig. 9, we repeat Fig. 7 with the error bars removed. Stars classed as metal-poor have been removed. Filled circles identify all stars with $[\text{Fe}/\text{H}]$ above -0.20 , triangles identify stars that are chromospherically active, open circles are stars with no definitive information, and the crosses show the location of stars within the Hyades as compiled in Twarog et al. (2002). In the range from $b - y = 0.50$ to 0.65 , the majority of the sample is tagged as metal-rich and occupies a position in the CMD consistent with that designation, *i.e.*, on or above the Hyades main sequence. Three stars bluer than $b - y = 0.66$ that occupy a position consistent with high $[\text{Fe}/\text{H}]$ but have no spectroscopic abundance determinations are HD 57095, HD 76378, and HD 219495.

Among the stars classed as chromospherically active, there is no apparent trend relative to the main sequence, with stars scattering above and below the Hyades, as expected if this sample originates from a population with a broader range in $[\text{Fe}/\text{H}]$. The majority of stars for which no information is available fall in the redder half of the distribution where spectroscopic data are few and far between; these stars scatter both above and below the Hyades. While it is possible that these stars represent a mixture of both metal-rich and chromospherically-active stars, we emphasize again the uncertainty in the definition of the mean Hyades relation among the coolest dwarfs. It may well be that with a larger array of cool Hyades and field star photometry, the position of the mean relation will shift or the samples for all metallicities will converge to a common relation insensitive to $[\text{Fe}/\text{H}]$, as originally expected for the K dwarfs.

5. Summary and Conclusions

An extensive sample of *Cuby* photometry of Hyades dwarfs from A through early M has been compiled and analyzed. For single stars and simple, composite, binary systems, the mean relation is well-defined over the color range from $b - y = 0.25$ to 0.65 . For the hotter portion of the color range, the effect of binarity on the two-color diagram is minimal. Among cooler dwarfs, the presence of a fainter, secondary star tends to shift the system above the mean relation, *i.e.*, to lower hk at a given $b - y$, simulating a lower $[\text{Fe}/\text{H}]$, with a maximum offset between 0.05 and 0.10 mag. Stars with extreme degrees of chromospheric activity, particularly *BY Dra* stars, appear anomalously metal-poor due to line-filling by emission within the Ca II H and K lines.

Among the F dwarfs, the Hyades relation defines a reliable lower bound to the distribution of field stars for a fixed $[\text{Fe}/\text{H}]$ of $+0.125$. By comparison to the spectroscopic metallicity scale of Edvardsson et al. (1993), it is found that the hk index is linearly correlated with $[\text{Fe}/\text{H}]$ over the range from $+0.5$ to -1.0 , with almost twice the sensitivity to abundance changes compared to m_1 . A simple test of the utility of hk at super-metal-rich levels is provided by the sample of stars recently identified as having planets as compiled by the University of California Planet Search Team (<http://exoplanets.org>). Of the 76 systems listed, 16 are in the hk Catalog (Twarog & Anthony-Twarog 1995). Of these, 6 are within the F-star color limits and have $[\text{Fe}/\text{H}]$ between $+0.17$ and $+0.34$, with a mean of $+0.23 \pm 0.08$. Line saturation does not appear to be a problem for the hotter stars.

In sharp contrast, as one extends the Hyades data toward cooler temperatures, the mean two-color relation crosses the distribution of nearby field stars, producing an approximate upper bound to the sample for stars in the color range from $b - y = 0.50$ to 0.65 . From an analysis of the stars that lie even higher than this relation in the $hk, b - y$ diagram, the primary source of the effect appears to be the high metallicity of the Hyades cluster. The implication is that the index does saturate at a given color for stars near solar abundance, but additional increases in $[\text{Fe}/\text{H}]$ lower the index, placing super-metal-rich stars in the same

region of the diagram as stars with $[\text{Fe}/\text{H}]$ near -0.7 or lower.

An alternative explanation may be provided by the synthetic indices developed from model atmospheres by Soon et al. (1993). While the models imply that hk should remain metallicity sensitive to $[\text{Fe}/\text{H}] = +0.5$ for all colors, a pattern almost identical to that found for hk is generated for C_{RV} , the photospheric narrow-band index constructed from the data collected for the Mt. Wilson survey of stellar chromospheric emission (Vaughan et al. 1978). Soon et al. (1993) explain the crossover of C_{RV} at cooler temperatures as primarily a result of the metallicity sensitivity of $b - y$, an effect that disappears with the use of $V - K$. Why this trend exists for an index based upon a filter over four times wider than those used in C_{RV} remains a mystery, but it may be more of an indication of the model shortcomings than of a serious problem with the observed indices. In any case, the hk index for cooler stars clearly succeeds in achieving the goal for which it was designed, the identification and calibration of stars of intermediate to extremely low $[\text{Fe}/\text{H}]$. This investigation adds two options to the list of uses, identification of stars with high levels of chromospheric activity, as expected, and the totally unexpected ability to isolate stars of Hyades abundance and higher. Three stars likely to occupy the last category and deserving of closer study are HD 57095, HD 76378, and HD 219495.

The question of the impact of modest, age-dependent variations in chromospheric emission remains unresolved, though indications from the very limited sample available to date are that the effect on hk is small to negligible. Observations of the G and K dwarfs in nearby open clusters of comparable age but lower $[\text{Fe}/\text{H}]$ than the Hyades could help to resolve this issue.

It is a pleasure to thank Dr. Rocha-Pinto for supplying us with a copy of the sample used in the study of the age-metallicity-relation from chromospheric emission. Although our thanks must reach back many years, we are indebted to the staffs of both NOAO observatories. We made extensive use of the WEBDA and SIMBAD databases, and are happy to acknowledge our appreciation to their creators and contributors. We are also indebted to the very helpful comments and constructive criti-

cisms of the referee.

REFERENCES

- Alexander, J. B. 1986, MNRAS, 220, 473
- Alonso, A., Arribas, S., & Martinez-Rogers, C. 1996, A&AS, 117, 227
- Anthony-Twarog, B. J., Laird, J. B., Payne, D., & Twarog, B. A. 1991, AJ, 101, 1902
- Anthony-Twarog, B. J., Sarajedini, A., Twarog, B. A., & Beers, T. C. 2000, AJ, 119, 2882
- Anthony-Twarog, B. J., Shawl, S. J., & Twarog, B. A. 1992, AJ, 104, 2229
- Anthony-Twarog, B. J., & Twarog, B. A., 1998, AJ, 116, 1902
- Anthony-Twarog, B. J., Twarog, B. A., & Craig, J. 1995, PASP, 107, 32
- Baird, S. R. 1996, AJ112, 2132
- Barry, D. C. 1974, AJ, 79, 616
- Bopp, B. W., Africano, J. L., & Goodrich, B. D. 1986, PASP, 98, 457
- Brown, J. A., 1983, AJ, 88, 678
- Carney, B. W. 1979, ApJ, 233, 211
- Cayrel de Strobel, G., Soubiran, C., & Ralite, N. 2001, A&A, 373, 159
- Chen, Y.Q. et al., 2000, A&AS141, 491
- Crawford, D. L. 1975, AJ, 80, 955
- Crawford, D. L., & Barnes, J. V. 1969, AJ, 74, 407
- Crawford, D. L. & Perry, C. L. 1966, AJ, 71, 206
- de Bruijne, J. H. J., Hoogerwerf, R., & de Zeeuw, P. T. 2001, A&A, 367, 111
- Deming, D., Olson, E. C., & Yoss, K. M. 1977, A&A, 57, 417
- Dobson, A. K. 1990, PASP, 102, 88
- Edvardsson, et al., 1993, A&AS, 275, 101
- Eggen, O. J. 1982, ApJS, 50, 22
- Eggen, O. J. 1994, AJ, 107, 594

- Favata, F., Micela, G., & Sciortino, S. 1997, *A&A*, 323, 809
- Griffin, R. F., Gunn, J. E., Zimmerman, B. A., & Griffin, R. E. M. 1988, *AJ*, 96, 172
- Griffin, R. F., & Gunn, J. E. 1981, *AJ*, 86, 588
- Grønbech, B., & Olsen, E. H. 1976, *A&AS*, 25, 213
- Hanson, R. B. & Vasilevskis, S. 1983, *AJ*, 88, 844
- Hauck, B., & Mermilliod, M. 1998, *A&AS*, 129, 431
- Hintz, M. L., Joner, M. D., & Hintz, E. G. 1998, *AJ*, 116, 2993
- Joner, M. D., & Taylor, B. J. 1995, *PASP*, 107, 124
- Laird, J. B., Carney, B. W., & Latham, D. W. 1988, *AJ*, 95, 1843
- Meusinger, H., Reimann, H. -C., & Stecklum, B. 1991, *A&A*, 245, 57
- Nissen, P. E. 1988, *A&A*, 199, 146
- Olsen, E. H. 1983, *A&AS*, 54, 55
- Olsen, E. H. 1993, *A&AS*, 102, 89
- Olsen, E. H. 1994, *A&AS*, 106, 257
- Patience, J., Ghez, A. M., Reid, I. N., Weinberger, A. J., & Matthews, K. 1998, *AJ*, 115, 1972
- Perry, C. L. 1969, *AJ*, 74, 705
- Perryman, M. A. C., Brown, A. G. A., Lebreton, Y., Gomez, A., Turon, C., Cayrel de Strobel, G., Mermilliod, J. -C., Robichon, N., Kovalevsky, J., & Crifo, F. 1998, *A&A*, 331, 81
- Reglero, V., Fabregat, J., Suso, J. 1992, *A&AS*, 92, 565
- Rey, S. -C., Lee, Y. -W., Joo, J. -M., Walker, A., & Baird, S. 2000, *AJ*, 119, 1824
- Rocha-Pinto, H. J., Maciel, W. J., Scalo, J., & Flynn, C. 2000, *A&A*, 358, 850
- Rose, J. A. 1984, *AJ*, 89, 1238
- Ryan, S. G., & Deliyannis, C. P. 1995, *ApJ*, 453, 819
- Schuster, W. J., & Nissen, P. E. 1988, *A&AS*, 73, 225
- Schuster, W. J., & Nissen, P. E. 1989, *A&A*, 221, 65
- Soderblom, D. R. 1989, *ApJ*, 342, 823
- Soon, W. H., Zhang, Q., Baliunas, S. L., Kurucz, R. L. 1993, *ApJ*, 416, 787
- Sowell, J.R. & Wilson, J. W. 1993, *PASP*, 105, 36
- Stauffer, J. R., Giampapa, M. S., Herbst, W., Vincent, J. M., Hartmann, L. W., & Stern, R. A. 1991, *ApJ*, 374, 142
- Strömgren, B., Gustafsson, B., & Olsen, E. H. 1982, *PASP*, 94, 5
- Taylor, B. J. & Joner, M. D. 1992, *PASP*, 104, 911
- Thorburn, J. A., Hobbs, L. M., Deliyannis, C. P., & Pinsonneault, M. H. 1993, *ApJ*, 415, 150
- Twarog, B. A. 1980, *ApJ*, 242, 242
- Twarog, B. A., & Anthony-Twarog, B. J. 1995, *AJ*, 109, 2828
- Twarog, B. A., & Anthony-Twarog, B. J. 1996, *AJ*, 112, 1500
- Twarog, B. A., Anthony-Twarog, B. J., & Tanner, D. 2002, *AJ*, submitted
- Twarog, B. A., Ashman, K. M., & Anthony-Twarog, B. A. 1997, *AJ*, 114, 2556
- Van Bueren, H. G. 1952, *Bull. Astr. Inst. Neth.*, 11, 385
- Vaughan, A. H. 1980, *PASP*, 92, 392
- Vaughan, A. H., Preston, G. W., & Wilson, O. C. 1978, *PASP*, 90, 267
- Wilson, O. C. 1963, *ApJ*, 138, 832

Fig. 1.— The effects of binarity on the $hk, b-y$ observations of cluster stars. (a) The CMD for stars combined with companions of decreasing temperature on the main sequence. (b) The $hk, b-y$ diagram for stars combined with companions of decreasing temperature on the main sequence.

Fig. 2.— The $hk, b-y$ diagram for the Hyades members. Open circles are single stars, crosses are probable multiple systems, and filled symbols are anomalous points excluded from the derivation of the mean relation, given by the solid curve.

Fig. 3.— $hk, b-y$ data for hotter field stars with the Hyades relation superposed.

Fig. 4.— Comparison of spectroscopic abundances from the catalogs of Edvardsson et al. (1993) and Chen et al. (2000), to the metallicity index, δhk . Solid line is a linear fit through the points.

Fig. 5.— Comparison of photometric abundances from the *uvby* photometry of Grønbech & Olsen (1976) (crosses) and Olsen (1983) (circles) to the metallicity index, δhk .

Fig. 6.— Same as Fig. 3 for the cooler field dwarfs.

Fig. 7.— CMD of stars located above the Hyades in the $hk, b-y$ diagram using *Hipparcos* parallaxes. Error bars show the uncertainty based upon the one sigma error in the parallax.

Fig. 8.— Abundance distribution for stars above the Hyades relation. Abundances represent a mixture of spectroscopic and photometric data. All stars with $[\text{Fe}/\text{H}]$ below -0.70 are contained in the lowest bin.

Fig. 9.— Same as Fig. 7 with evolved stars and metal-poor stars removed. Filled circles are the stars with $[\text{Fe}/\text{H}]$ above -0.2 , triangles are stars that are chromospherically active, open stars are data points without any clarifying information, and crosses are observed single stars within the Hyades.